



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C07C 61/06</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/28255</b> <b>(43) International Publication Date:</b> 2 July 1998 (02.07.98)
<b>(21) International Application Number:</b> PCT/US97/23164 <b>(22) International Filing Date:</b> 24 December 1997 (24.12.97) <b>(30) Priority Data:</b> 119890 24 December 1996 (24.12.96) IL <b>(71) Applicant (for all designated States except BB US):</b> TEVA PHARMACEUTICAL INDUSTRIES LTD. [IL/IL]; Basel Street 5, P.O. Box 3190, 49131 Petah Tiqva (IL). <b>(71) Applicant (for BB only):</b> TEVA PHARMACEUTICALS USA, INC. [US/US]; 1510 Delp Drive, Kulpsville, PA 19443 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> PESACHOVICH, Michael [-/IL]; Emek Dotan 16/7, Ramat Gan (IL). SINGER, Claude [-/IL]; David Elazar 8/8, Kfar Sava (IL). PILARSKI, Gideon [-/IL]; Ataroth 12/29, Holon (IL). <b>(74) Agents:</b> LEE, Steven, J. et al.; Kenyon & Kenyon, One Broadway, New York, NY 10004 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> PREPARATION OF GABAPENTIN  <b>(57) Abstract</b>  There is disclosed a method of converting gabapentin hydrochloride substantially free of inorganic salts to gabapentin form II. The method comprises the steps of: (1) obtaining gabapentin hydrochloride that is substantially free of inorganic salts; (2) mixing a solution of the gabapentin hydrochloride with an additional amine in a first solvent so as to obtain a precipitate comprising gabapentin; and then (3) recovering gabapentin form II from the precipitate. The precipitated gabapentin is a novel polymorphic form of gabapentin possessing a crystalline structure characterized by novel sets of peaks in the powder X-ray diffraction pattern and in the FTIR spectra. This newly disclosed polymorph of gabapentin is characterized herein as gabapentin form III. The recovery step may comprise, for example, one of two alternative methods, slurring the gabapentin form II in methanol, and then filtering the suspension to obtain gabapentin form II, or solubilizing the gabapentin form III in methanol with heating by reflux, and then cooling the solution to obtain gabapentin form II by crystallization.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		


PREPARATION OF GABAPENTIN

5    FIELD OF THE INVENTION

This invention relates to a new process for converting gabapentin hydrochloride salt to gabapentin via a novel polymorphic form of gabapentin.

10   BACKGROUND OF THE INVENTION

Gabapentin is 1-(aminomethyl)-1-cyclohexaneacetic acid, having the chemical structure:



- 15   Gabapentin is used in the treatment of cerebral diseases such as epilepsy. The literature describes many ways of preparing gabapentin from a variety of starting materials. U.S. Patent 4,024,175 describes at least three methods of preparing gabapentin from cyclohexyl-1,1-diacetic acid. Each of these
- 20   methods results in the formation of gabapentin hydrochloride salt, which may be converted to 1-(aminomethyl)-1-cyclohexaneacetic acid by treatment with a basic ion exchanger and then crystallized from a solvent such as ethanol/ether.
- 25   U.S. Patent 4,894,476 specifically discloses an improved method for converting the hydrochloride salt into the free amino acid. This involves pouring a deionized water solution of the salt over an ion exchange column, eluting with deionized water, producing a slurry from the eluate, adding an

alcohol to the slurry, centrifuging and drying the slurry to obtain the free amino acid.

Alternative methods for preparing gabapentin have been  
5 described that do not proceed via the hydrochloride or any  
other mineral acid salt. Such methods include those described  
in U.S. Patent Nos. 5,132,451, 5,095,148, 5,068,413. Each of  
these methods involve a cyanic intermediate which is  
hydrogenated under severe conditions to produce the free amino  
10 acid.

These methods are industrially impractical. Those methods  
comprising ion exchange columns require the use of large  
amounts of ion exchanger for lengthy periods of time to lower  
15 the level of chloride ions to the desired level. The  
alternative methods involve further more demanding steps.

Commercially available gabapentin is crystalline and exhibits  
an X-ray diffraction pattern with peaks of 2-theta values at  
20 7.8, 13.3, 15.0, 17.0, 20.4, 21.3, 23.1, 23.6, 25.7, 27.0 and  
28.2 degrees. Hereinafter, the commercially available  
polymorphic form of gabapentin is referred to as polymorph  
form "II".

## 25 SUMMARY OF THE INVENTION

The present invention relates to an improved method for  
purifying gabapentin comprising converting gabapentin  
hydrochloride salt to gabapentin form II. The present  
invention avoids the disadvantages associated with prior art  
30 methods, by adding alternative steps and by proceeding via a  
novel polymorphic form of gabapentin.

Accordingly, the present invention relates to a method of  
converting gabapentin hydrochloride salt to gabapentin form  
35 II, comprising reacting a solution of gabapentin hydrochloride

with an additional amine in a first solvent to produce a novel polymorphic form as a precipitate, thereafter, converting the novel polymorphic form to form II by forming a suspension and/or a solution of the precipitate in methanol, and then  
5 recovering gabapentin form II.

The present invention further relates to a novel polymorphic form of gabapentin designated as gabapentin form III. The polymorph may be identified by its unique X-ray diffraction  
10 pattern.

A further aspect of the present invention relates to the novel polymorphic form of gabapentin that is of use as an intermediate in the preparation of polymorphic form II.  
15

Further objectives and advantages of the subject invention will be apparent to those skilled in the art from the detailed description of the disclosed invention.

## 20 **BRIEF DESCRIPTION OF THE DRAWINGS**

Fig. 1 is an x-ray diffraction pattern of a sample of gabapentin hydrate.

Fig. 2 is an x-ray diffraction pattern of gabapentin form II.  
25

Figs. 3a and 3b are x-ray diffraction pattern of gabapentin form III, possibly containing small amounts of gabapentin form II and/or gabapentin hydrate.

30 Fig. 4 is an FTIR (Fourier Transform Infra Red) spectrum of gabapentin hydrate.

Fig. 5 is an FTIR spectrum of gabapentin form II.

35 Fig. 6 is an FTIR spectrum of gabapentin form III, possibly

containing small amounts of gabapentin form II and/or gabapentin hydrate.

**DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS**

5 The subject invention will now be described in detail for specific preferred embodiments of the invention, it being understood that these embodiments are intended only as illustrative examples and the invention is not to be limited thereto.

10

The method of the present invention comprises:

- (1) obtaining gabapentin hydrochloride that is substantially free of inorganic salts;
- (2) mixing a solution of the gabapentin hydrochloride  
15 with an additional amine in a first solvent so as to obtain a precipitate comprising gabapentin; and then
- (3) recovering gabapentin form II from the precipitate.

Preferably, the gabapentin hydrochloride used as a starting  
20 material in the process of the present invention is substantially free of other inorganic salts such as sodium chloride and sodium bromide, that is, such impurities are only present in trace amounts. Alternatively, gabapentin hydrochloride containing inorganic salts may be used subject  
25 to the addition of a further step for removing the inorganic salts prior to mixing gabapentin hydrochloride with the additional amine.

Thus, gabapentin hydrochloride containing inorganic salts may  
30 optionally be pre-treated to remove the inorganic salts by the steps of (a) dissolution in a solvent in which gabapentin hydrochloride is soluble, but the inorganic salts are not; (b) filtration of the inorganic salts and, optionally, (c) evaporating the solvent to recover gabapentin hydrochloride  
35 substantially free of inorganic salts. Appropriate solvents

for the optional pre-treatment step include those selected from the group consisting of methanol, ethanol, n-propanol, isopropanol, butanol, t-butanol, n-butanol, ethyleneglycolmonomethylether, benzylalcohol or  
5 dimethylacetamide. Preferably the pre-treatment solvent is identical to the first solvent. This pre-treatment solvent, which is identified in Table 1 and in the claims as the "second solvent", would be used prior to using the "first solvent", if, in fact, this optional pre-treatment step is  
10 included in the process.

The second step of the present invention comprises mixing a solution of the gabapentin hydrochloride in a solvent that contains an additional amine. The solvent may be any solvent  
15 in which the hydrochloride salt of the additional amine is soluble but in which gabapentin form III is insoluble, such that a precipitate of gabapentin form III is formed. Such solvents are preferably selected from the group consisting of ethyl acetate, dimethylcarbonate, ethanol, butanol, t-butanol,  
20 n-butanol, methanol, acetonitrile, toluene, isopropylacetate, isopropanol, methylethylketone, acetone, ethyleneglycolmonomethylether, methylene chloride, chloroform, benzylalcohol or dimethylacetamide. The precipitated gabapentin, which may be separated by filtration, is  
25 characterized herein as a novel polymorphic form of gabapentin, possessing a crystalline structure characterized by peaks in the powder X ray diffraction pattern with 2-theta values at 6.11, 12.22, 17.00, 18.20, 19.94, 20.81, 24.54, and 25.11 degrees, all  $\pm 0.2$  degrees. The x-ray diffraction  
30 pattern in the samples obtained appears to show a preferred orientation in which the peak at 6.11 degrees is larger than any other peak in the pattern, and the peaks at 12.22 and 24.54 degrees are larger than any of the remaining peaks in the pattern. This polymorph is referred to herein as  
35 gabapentin form III.

Two representative embodiments of the third step of the present invention, recovery of gabapentin form II from the precipitate, are slurring and crystallization. Slurring may be performed by suspending the precipitated gabapentin in  
5 methanol by mixing, stirring, and/or providing continuous agitation with some mechanical device so as to induce transformation into the gabapentin form II that is the commercially available polymorphic form of gabapentin. Gabapentin form II may then be filtered off and washed.

10

Alternatively, the precipitated gabapentin may be crystallized from methanol with heating by reflux until dissolved, cooling, optionally seeding with gabapentin, followed by further cooling, and then collecting and drying the crystals of  
15 gabapentin form II. Second and multiple crops may be obtained from the concentrated mother liquors.

Suitable amines for use in the present invention include triethylamine, tributylamine, tripropylamine, trihexylamine,  
20 diethylamine, ethanolamine and benzylamine. Preferably the amine is tributylamine.

The form II gabapentin obtained by the methods of the present invention may be crystallized using processes known in the  
25 art.

Certain specific representative embodiments of the invention are described in detail below, the materials, apparatus and process steps being understood as examples that are intended  
30 to be exemplary and illustrative only. In particular, the invention is not intended to be limited to the methods, materials, conditions, process parameters, apparatus and the like specifically recited herein.

35 **EXAMPLES**



# 1. Preparation of gabapentin from gabapentin hydrochloride

## A. Removal of Inorganic Salts

Gabapentin hydrochloride containing inorganic salts (NaCl, NaBr), which may be prepared, for example, as shown in U.S. Pat. No. 4,024,175, Example 4, Variant A, was dissolved in 150 ml isopropanol and mixed at 25°C. Active carbon (0.9 g) was added and the suspension mixed for a further 2 hours. The inorganic salts were removed by filtering. The filter cake was washed twice with 15 ml isopropanol and the washings were added to the gabapentin hydrochloride solution.

## B. Precipitation of Gabapentin Base; Formation of Gabapentin Form III

The gabapentin hydrochloride solution of part A was concentrated to dryness in vacuum while ensuring that the temperature of the heating bath did not exceed 35°C. 210 ml ethylacetate and 16.5 ml tributylamine were added and the solution was mixed for 2 hours at 25°C. The gabapentin precipitate was then separated by filtration, the filter cake being washed with 20 ml ethylacetate and then 20 ml methanol. The filter cake contained gabapentin form III which when dried displayed a characteristic X-ray diffraction pattern with 2-theta values at 6.1, 12.2, 17.0, 17.7, 18.3, 20.0, 20.8, 24.6 and 25.5 degrees.

## C. Alternative Methods for Conversion of Precipitated Gabapentin to Gabapentin Form II

### (1) Slurrying

The still humid filter cake from step 2 was suspended in 52.5 ml methanol for about 14 hours at 25°C. Solid gabapentin was then separated from the suspension by filtration. The filter cake was washed with 20 ml methanol and then dried under vacuum at 35°C. 10.8 g crystalline gabapentin form II (yield 72%) was obtained.

(2) Crystallization

The still humid filter cake from step 2 was suspended in 210 ml of methanol and the suspension was heated to reflux (65°C). If dissolution was incomplete, additional methanol was added.

After dissolution, the reactor content was cooled to 34°C at which temperature crystallization was induced by seeding with pure Form II gabapentin base (0.1 g).

After maintaining the mixture at 34°C for 60 minutes the reactor content was cooled to 25°C and methanol was distilled by vacuum distillation. Approximately 160 ml of methanol was collected. Then the suspension was cooled to 0-10°C and maintained at this temperature for 2 hours.

The crystalline gabapentin was separated by filtration from the suspension. The filter cake was washed with 20 ml methanol and then dried under vacuum at 35°C. 10.8 g of crystalline gabapentin form II (yield 72%) was thus obtained.

Examples 2-20

The method of Example 1 was followed using the slurring technique of step C(1) and employing the amines and solvents shown in Table 1 below. The percent yields are of purified product except where marked with an asterisk (\*), where the yield was measured at the stage prior to the slurring in methanol.

Table 1. Summary of the first solvent, second solvent, amine and yield of examples 2-20.

	Example No.	First Solvent	Second Solvent	Amine	Yield(%)
5	2	IPA	IPA	TBA	61.5
	3	Et Ac	IPA	TBA	74.0
	4	Et Ac	IPA	THA	66.2
	5	Et Ac	IPA	TPA	63.0
	6	Et Ac	IPA	THA	79.6*
10	7	IPA Ac	IPA	TBA	64.9
	8	ACN	IPA	TBA	67.8
	9	DMC	IPA	TBA	57.9
	10	DMA	DMA	TBA	65.7
	11	BzOH	BzOH	TBA	43.0*
15	12	MEK	IPA	TBA	88.0*
	13	t-BuOH	t-BuOH	TBA	79.4*
	14	Acetone	IPA	TBA	73.1*
	15	Et Ac	BuOH	TBA	69.8
	16	MeOH	MeOH	TBA	67.4
20	17	EGMME	EGMME	TBA	66.8*
	18	IPA	IPA	TEA	76.2*
	19	IPA	IPA	BzA	56.0*
	20	CH <sub>2</sub> Cl <sub>2</sub>	DMA	DEA	89.4*

- 25 IPA isopropanol                      MEK methyl ethyl ketone  
 ACN acetonitrile                      EGMME ethyleneglycol monomethylether  
 BzOH benzyl alcohol                      TBA tributylamine  
 MeOH methanol                      BzA benzylamine  
 CH<sub>2</sub>Cl<sub>2</sub> methylene chloride                      IPA Ac isopropylacetate  
 30 THA trihexylamine                      DMC dimethylcarbonate  
 TPA tripropylamine                      (t)-BuOH (tert)-butanol  
 Et Ac ethyl acetate                      TEA triethylamine

DMA dimethylacetamide

DEA diethylamine

The gabapentin form III was characterized by comparing the x-ray diffraction pattern and the FTIR absorption spectra of gabapentin form III with gabapentin form II and gabapentin hydrate. The x-ray powder diffraction patterns of Figures 1, 2, and 3a were obtained using a Philips x-ray powder diffractometer with the following parameters:

10 Goniometer model 1050/70, Cu-tube, Curved graphite monochromator.

Sample holder: Quartz monocrystal plate

Settings:

X-Ray tube:

15 KV-40

mA-28

Target-Cu

Divergence slit-1°

Receiving parallel slit-0.2 mm

20 Scatter slit-1°

Scintillation detector:

Voltage-832

Lower level-33.8%

Window-34.5%

25 Scan parameters:

Scanning speed: 2°/min

Paper speed: 2cm/min

Gain: 32

30 Calibration: External calibration with silicon fine powder

Type of radiation: copper K $\alpha$ .

Figure 3b was obtained using a Siemens B5100 with a presample (K $\alpha_1$  only) monochromator, step scan, Cu radiation, beam slits 0.3, receiving slit 0.05, standard Siemens rotating sample

holder, start/stop/step angles in degrees were 4.0, 35, and 0.04, with a step duration of 15 seconds. Standard Siemens D-5000 software was used. The diffractometer was calibrated with NIST large d-spacing standard. No theta-compensating slits were employed.

The x-ray spectra of gabapentin hydrate, gabapentin form II and gabapentin form III are shown in Figs. 1-3, respectively. The main diffraction peaks that characterize each material are listed in Table 2.

Table 2. X-ray Diffraction peaks of gabapentin hydrate, gabapentin form II and gabapentin form III.

Hydrate	Form II	Form III
2-theta (°)	2-theta (°)	2-theta (°)
6.1	7.9	6.11
12.2	13.3	12.22
16.0	15.0	17.00
18.3	17.0	17.63
19.1	19.5	18.20
19.8	20.3	19.94
20.7	21.3	20.81
24.5	21.8	24.54
26.4	23.0	25.11
28.4	23.6	28.91
30.7	25.7	30.20
32.3	26.9	30.78
	28.2	31.46

The FTIR spectra for gabapentin hydrate, gabapentin form II and gabapentin form III are shown in Figs. 4-6,

respectively. The FTIR peaks are summarized in Table 3.

Table 3. FTIR peaks of gabapentin hydrate, gabapentin form II and gabapentin form III.

5

10

15

20

Form I	Form II	Form III
cm <sup>-1</sup>	cm <sup>-1</sup>	cm <sup>-1</sup>
648	709	708
726	749	760
880	890	885
926	922, 928	926
968	976	974
1154	1165	1160
1175		1180
1292	1300	1290
	1420	1420
	1476	1460
		1510
1542	1546	1586
1624	1615	1664
1662		

Morphology of the three forms:

- 25 1. The hydrate form typically exists as large crystals with undefined shapes.
  2. Form II typically exists as plate shaped crystals.
  3. Form III typically exists as small rhomboidal crystals.
- 30 The melting point for gabapentin was not determined since gabapentin decomposes prior to melting.

**What Is Claimed Is:**

1. A method of converting gabapentin hydrochloride to gabapentin form II comprising:
  - (1) dissolving gabapentin hydrochloride in a first solvent in which gabapentin is relatively insoluble; and
  - (2) adding an additional amine to the gabapentin hydrochloride solution so as to obtain a precipitate comprising gabapentin.
2. The method of claim 1, further comprising the step of recovering gabapentin form II from said precipitate.
3. The method of claim 2, the recovering step comprising slurrying or crystallizing said precipitate in methanol.
4. The method of claim 3, wherein said first solvent is selected from the group consisting of ethyl acetate, dimethylcarbonate, ethanol, butanol, t-butanol, n-butanol, methanol, acetonitrile, toluene, isopropylacetate, isopropanol, methylethylketone, acetone, ethyleneglycolmonomethylether, methylene chloride, chloroform, benzylalcohol or dimethylacetamide.
5. The method of claim 1 wherein said additional amine is selected from the group consisting of triethylamine, tributylamine, tripropylamine, trihexylamine, diethylamine, ethanolamine and benzylamine.
6. The method of claim 5 wherein said additional amine is tributylamine.
7. The method of claim 1 wherein said first solvent is selected from the group consisting of ethyl acetate, dimethylcarbonate, ethanol, butanol, t-butanol, n-butanol, methanol, acetonitrile, toluene, isopropylacetate,

isopropanol, methylethylketone, acetone, ethyleneglycolmonomethylether, methylene chloride, chloroform, benzylalcohol or dimethylacetamide.

8. The method of claim 7 wherein said first solvent is ethylacetate.

9. The method of claim 1 wherein said first solvent is one in which the hydrochloride salt of said additional amine is soluble.

10. The method of claim 1 further comprising pre-treating said gabapentin hydrochloride to remove inorganic salts, which pre-treatment comprises (a) dissolving gabapentin hydrochloride in a second solvent in which inorganic salts are insoluble, (b) filtering off the inorganic salts, and (c) recovering gabapentin hydrochloride from said second solvent.

11. The method of claim 10 wherein said second solvent is selected from the group consisting of methanol, ethanol, n-propanol, isopropanol, butanol, t-butanol, n-butanol, ethyleneglycolmonomethylether, benzylalcohol or dimethylacetamide.

12. The method of claim 11 wherein said second solvent is isopropanol.

13. The method of claim 1 or 10 wherein the precipitated gabapentin comprises gabapentin form III.

14. The method of claim 10 wherein the precipitated gabapentin comprises gabapentin form III, said second solvent is isopropanol, said first solvent is ethylacetate, and said additional amine is tributylamine.



15. The method of claim 14 wherein said precipitated gabapentin is slurried or crystallized from methanol to yield gabapentin form II.

16. Crystalline gabapentin form III characterized by powder X-ray diffraction peaks at  $6.11 \pm 0.2$ ,  $12.22 \pm 0.2$ ,  $17.00 \pm 0.2$ ,  $17.63 \pm 0.2$ ,  $18.20 \pm 0.2$ ,  $19.94 \pm 0.2$ ,  $20.81 \pm 0.2$ ,  $24.54 \pm 0.2$ ,  $25.11 \pm 0.2$ ,  $28.91 \pm 0.2$ ,  $30.20 \pm 0.2$ ,  $30.78 \pm 0.2$ , and  $31.46 \pm 0.2$  degrees 2-theta.

17. Crystalline gabapentin form III characterized by powder X-ray diffraction peaks at  $6.11 \pm 0.2$ ,  $12.22 \pm 0.2$ ,  $17.00 \pm 0.2$ ,  $17.63 \pm 0.2$ ,  $18.20 \pm 0.2$ ,  $19.94 \pm 0.2$ ,  $20.81 \pm 0.2$ ,  $24.54 \pm 0.2$ ,  $25.11 \pm 0.2$ ,  $28.91 \pm 0.2$ ,  $30.20 \pm 0.2$ ,  $30.78 \pm 0.2$ , and  $31.46 \pm 0.2$  degrees 2-theta, wherein the size of the peaks is in the order  $6.11$  degrees  $> 12.22$  and  $24.54 > 17.00$ ,  $17.63$ ,  $18.20$ ,  $19.94$ ,  $20.81$ ,  $25.11$ ,  $28.91$ ,  $30.20$ ,  $30.78$ , and  $31.46$ .

18. Crystalline gabapentin form III characterized by infra-red absorptions having peaks at 708, 760, 885, 926, 974, 1160, 1180, 1290, 1420, 1460, 1510, 1586, and  $1664 \text{ cm}^{-1}$ .

19. Crystalline gabapentin form III produceable by a process comprising the steps of:

(a) dissolving gabapentin hydrochloride in a solvent selected from the group consisting of ethyl acetate, dimethylcarbonate, ethanol, butanol, t-butanol, n-butanol, methanol, acetonitrile, toluene, isopropylacetate, isopropanol, methylethylketone, acetone, ethyleneglycolmonomethylether, methylene chloride, chloroform, benzylalcohol or dimethylacetamide;

(b) adding an additional amine selected from the group consisting of triethylamine, tributylamine, tripropylamine, trihexylamine, diethylamine, ethanolamine and benzylamine; and

(c) drying the precipitated gabapentin.

Figure 1




Figure 2








Figure 3b



5/7


Figure 4



95/11/07 18:15 ZIONA TEVA AL-IR-40  
m071195b: 16 scans, 4.0cm-1  
GABAPENTIN: gabapentine monohydrate. KBr disk

6/7


Figure 5



97/01/15 15:01 ZIONA TEVA AL-IR-40  
m150197a: 16 scans, 4.0cm-1  
GBP GP-1511, 1% in KBr form II

7/7

Figure 6



96/06/27 18:28 ZIONA TEVA AL-IR-40  
m270696a: 16 scans, 4.0cm-1  
GABAPENTIN AL-9244-3(KBr) form III



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/23164

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07C 61/06

US CL : 562/507

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 562/507

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- A	US 4,024,175 A (SATZINGER et al.) 17 May 1977, col. 4, lines 22-24.	16-19 ----- 1-15
X -- A	US 4,960,931 A (BUTLER et al.) 02 October 1990, Example 3.	16-19 ----- 1-15
X -- A	US 5,068,413 A (STEINER et al.) 26 November 1991, Examples 3 and 6.	16-19 ----- 1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

20 MARCH 1998

Date of mailing of the international search report

21 APR 1998

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MICHAEL L. SHIPPEN

Telephone No. (703) 308-1235